

Stature estimation from facial measurements in medical students of Tehran university of Medical Sciences: an Iranian population

Fatemeh Navaei, Neda Ghaffari, Sina Mojaverrostami, Masoumeh Dodongeh, Mostafa Nemati and Gholamreza Hassanzadeh

Department of Anatomy, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran.

Correspondence to Gholamreza Hassanzadeh (email: hassanzadeh@tums.ac.ir).

(Submitted: 27 April 2018 – Revised version received: 04 June 2018 – Accepted: 14 July 2018 – Published online: 26 September 2018)

Objectives Stature (*S*) estimation is one of the most important methods for identification of individual in forensic medicine. Different kinds of methods are available for estimation of *S* from different parts of the body.

Methods The present study aimed to estimate *S* from three facial measurements including facial breadth (FB), total face height (TFH) and upper facial height (UFH) of 200 healthy students (100 males and 100 females) aged 20–25 years, studying at Tehran University of Medical Sciences. The correlation between stature and facial anthropometric measurements was investigated and equations for stature prediction were demonstrated by using linear regression analysis.

Results The mean stature, FB, TFH and UFH in males were higher than females. As well, there were significant correlations between stature and FB, TFH and UFH in all subjects, males and females. Additionally, we could formulate the stature from FB, TFH and UFH by using linear regression equation.

Conclusion According to the results, the formulas were calculated for predicting stature from facial measurements. So in forensic and archeology investigations, if the available remains are skulls, to predict the stature we can use the obtained formulas from Iranian population.

Keywords stature, facial measurements, anthropometry, regression

Introduction

Anthropometric studies are essential in forensic medicine and archeology for identifying a true race from the local mingling of races.¹ Stature is one of the most frequently used biological profiles for identification of an individual.^{2,3} One of the main goals of forensic medicine is to identify an unknown or tentative person, particularly when just some parts of deceased body remain.⁴ The stature is the height of a person in standing position, which is varied in different ethnic groups and populations.⁵ This variation in stature measurement of different populations is related to the several factors such as genetic, nutritional status and environmental factors.⁶ The relationship of measurements between the different body segments and the stature were analyzed in different countries and equations for stature estimation were determined.^{7–9} Stature and sex can be predicted from different body measurements by using statistical models.¹⁰ The improvement of models to predict sex and stature from anthropometric data in different ethnic groups are necessary, because there is a great need for population-specific studies on stature estimations.¹¹ Several studies have been performed on the estimation of stature from different body parts such as hand length,¹² foot length¹³ and metacarpal length.⁶

The craniofacial anthropometry is a branch of anthropometry which deals with head and face measurements.¹⁴ Craniofacial morphology has great importance in maxillofacial surgery, plastic surgery, orthodontics and forensic medicine.¹⁵ Craniofacial measurements are varied from one person to another due to several influencing factors such as age, sex, ethnicity, nutrition, genetic, environmental conditions and soft-tissue matrices.¹⁶ Sometimes from an unknown fragmentary deceased just the skull is available. In this situation, cephalo-facial dimensions are very useful for accurate identification.¹⁷ Different facial measurements were used for stature estimation as a decisive factor for identifying an

individual such as total and upper facial height, height of lower face, minimum frontal, bigonial, biocular and interocular breadth.¹⁸

This study aimed to evaluate the correlation between three facial measurements including facial breadth (FB), total facial height (TFH) and upper facial height (UFH) and stature in medical students of Tehran University of Medical Sciences.

Materials and Methods

Subjects

This study was designed to evaluate among the students of Tehran University of Medical Sciences in Iran. A total number of 200 students were randomly selected from the volunteers of Iranian students (100 females and 100 males) in the age range of 20–15 years old with normal craniofacial configuration, without any deformity of the vertebral column and no history of facial plastic surgery. Each subject was examined for stature, FB, TFH and UFH. Afterward, the association between stature and mentioned parameters was determined by statistical analysis.

Stature

Stature was measured by a standard anthropometer in centimeters. The student was asked to stand barefoot on the flat surface base in an erect position with the head oriented in the Frankfurt Plane. Then, the measurement was taken in centimeters as the vertical distance between the vertex to the flat surface.¹⁹

Facial measurements

Facial measurements were taken by using sliding venire calipers.

Facial breadth

Measured as a distance between two zygions (Zy).

Total facial height

Measured as a distance between nasion (N) and menton (Me) in centimeters.

Upper facial height

Measured as a distance between N and subnasale (Sn) in centimeters (20).

Definition

Zy: Most lateral point of zygomatic arches.

N: Midline depression locating below the level of the eyebrows.

Sn: Point of the nasal septum and the upper lip Junction.

Me: Inferior point on the symphysis (the lowest portion of chin).

Statistical analysis

The data analysis was performed using SPSS 22.0 for Windows. The differences between two sexes were tested by using independent *t*-test. The association between stature and facial measurements was determined by using Pearson correlation analysis. Linear regression analysis was used to determine equations for stature estimation from facial measurements. $P < 0.05$ was considered significant.

Results

According to Table 1, there were significant differences in stature ($P = 0.0001$) and facial anthropometric measurements including FB ($P = 0.02$), TFH ($P = 0.0001$) and UFH ($P = 0.0001$) of sex groups.

The correlation confections between stature and facial anthropometric measure were evaluated in Table 2. There were signification correlation confections between S and FB ($r = 0.002$ and $P = 0.001$), TFH ($r = 0.02$ and $P = 0.0001$) and UFH ($r = 0.01$ and $P = 0.0001$). The male and females were analyzed separately. As shown in Table 2, the significant correlation were recorded for male subjects between S and FB ($r = 0.001$ and $P = 0.0001$), TFH ($r = 0.01$ and $P = 0.0001$) and UFH ($r = 0.01$ and $P = 0.0001$). As well, these values were lower for female subjects as defined between S and FB ($r = 0.0001$ and $P = 0.003$), TFH ($r = 0.001$ and $P = 0.002$) and UFH ($r = 0.01$ and $P = 0.01$).

Based on Table 3, the linear regression was performed to find reliable formulas between S and FB, TFH and UFH. Additionally, the relation between S and FB, TFH and UFH were formulated for male and female subjects, separately.

Discussion

In the forensic field, stature is an important and reliable factor for personal identification in addition to age, sex, and ethnical origin and is a factor that helps forensic scientist to narrow the medicolegal investigations. According to literature, formulation of stature by using long bones of limbs is one of the most accurate factors.^{21,22} However, when the long bones are not available and remains of skull, head or face are found in the forensic investigations, it becomes hard for forensic specialist to identify the cases. So characteristics of other bones or body parts such as skull or head and neck have been performed to predict stature.¹⁸ In different studies, it has been demonstrated that regression analysis is the most reliable and easy method for predicting the stature.^{23,24} In this study, we focused to formulate the stature from face measurements by using regression analysis in an Iranian population, rarely studied.

At first, we compared the measurements based on sex groups. Our results showed that the facial measurements were higher than females. Different studies confirmed our results. Din et al.²⁵ showed that the nasofacial values in females were lower than males. In a study by Omotoso et al.,²⁶ they demonstrated that the mean nasofacial measurements were different based on the gender. Their results are comparable to the present study.

Studied on stature prediction based on cephalo-facial anthropometric characteristic are so limited. We could estimate correlation confections between stature and facial measurement for total studies sample and separately for male and female subjects. The strongest one was between TFH and stature. In a study on Koli male subjects by Krishan,²⁷ the high correlation between cephalo-facial dimensions and stature was reported for North India population. Their findings were related to the Koli population as an endogamous group with high homogeneity.²⁷ As well, in a study on Japanese cadavers, Chiba and Terazawa²⁸ reported a correlation between cephalo-facial values and stature with correlation coefficients range of 0.32–0.53. After excluding over 70 year subjects, they reported higher correlation coefficients between stature and these measurements.²⁸

Additionally, our results confirmed that the male subjects had stronger correlation coefficient compared to the female subjects. Chiba and Terazawa²⁸ recorded that female subjects had shown smaller correlation coefficient than males. Chiba and Terazawa²⁸ also emphasized that the morphological structure of the head, especially the antero-posterior diameter, changed in relation to the age of the individual. Kalia et al.²⁹ tried to estimate stature by using

Table 1. Comparing facial anthropometric measurements based on sex groups *n* medical students of Tehran University of Medical Sciences

	Total				Sex								P-value
					Male				Female				
	M	SD	Min	Max	M	SD	Min	Max	M	SD	Min	Max	
FB (cm)	11	0.11	8	13	11.2	0.1	10	13	12	0.2	8	112	0.02
TFH (cm)	11	0.12	8	13	11.1	0.1	9	13	10	0.1	8	11	0.0001
UFH (cm)	7.05	0.2	5	8	7.1	0.2	6	8	6	0.2	5	8	0.0001
S (cm)	171	9.3	153	190	178.2	6.1	163	194	164	5.02	153	175	0.0001

M, mean; SD, standard deviation; min, minimum; max, maximum; FB, facial breadth; TFH, total facial height; UFH, upper facial height; S, stature.

Table 2. The correlation between stature and facial anthropometric measurements

		FB (cm)	TFH (cm)	UFH (cm)
S_f (cm)	<i>r</i>	0.002	0.02	0.01
	<i>P</i> -value	0.001	0.0001	0.0001
S_m (cm)	<i>r</i>	0.001	0.01	0.01
	<i>P</i> -value	0.0001	0.0001	0.0001
S_f (cm)	<i>r</i>	0.0001	0.001	0.01
	<i>P</i> -value	0.003	0.002	0.01

r, correlation coefficients; FB, facial breadth; TFH, total facial height; UFH, upper facial height; *S*, stature; m, males; f, females.

Table 3. Formulation of stature by using facial measurements in total, males and females

Formulas	R ²	SEM	<i>P</i> -value
$S_f = 3.014FB + 136$	0.051	6	0.0001
$S_f = 6TFH + 98$	0.062	6	0.0001
$S_f = 8UFH + 115$	0.041	7	0.0001
$S_m = FB + 160$	0.043	6	0.0001
$S_m = 3TFH + 137$	0.061	6.5	0.0001
$S_m = 8UFH + 115$	0.066	6	0.0001
$S_f = FB + 149$	0.044	4	0.0001
$S_f = TFH + 148$	0.037	4	0.0001
$S_f = 2UFH + 140$	0.055	4.2	0.0001

SEM, standard error of the estimate; FB, facial breadth; TFH, total facial height; UFH, upper facial height; *S*, stature; m, males; f, females

odontometry and skull anthropometry. Correlation coefficients were between 0.56 and 0.38 and were close to those

of Chiba and Terazawa, but only when combined data were used. When male and female individuals were evaluated separately, a significant decrease was observed in the correlation coefficients. For male subjects, they changed between 0.20 and 0.13, which is similar to the results of this study. However, Sarangi et al.³⁰ investigated the correlations between stature and maximum transverse length circumference of the skull and recorded that the correlation coefficients of stature for those variables were insignificant for a suitable estimation of stature.

In our study, the range of SEE was 6–7 for all subjects, 6–6.5 for male and 4–4.5 for females according to the findings obtained from linear regression equations. *R*-square obtained from regression equation for facial measurements was less than 0.1. In the prediction of *S* from facial measurements in northwest Indians, Sahni et al.¹⁸ demonstrated that the range of SEE obtained from multiple regression was 3.569–3.610 in males and 2.880–2.914 in females. Agnihotri et al.³¹ used cephalo-facial dimensions to estimate stature by regression analysis in Indo-Mauritian population. Their results showed that *r* of all cephalo-facial measurements were less than 0.5. All these findings confirm that the correlation coefficients are not high for facial measurement to predict stature and the findings are directly related to the homogeneity of populations studied.

Conclusion

According to the findings of the present study, the formulas were calculated from facial measurements for predicting stature in an Iranian population. As well, the correlation coefficients calculated for males were better than females. Forensic scientists and anthropologists can use from these findings to identify cases which just their remains are head, face or skull in Iranian population. ■

References

- Mohammed I, Mokhtari T, Ijaz S, Ngaski AA, Akanji Omotosho D, Milanifard M, et al. Anthropometric study of nasal index in Hausa ethnic population of northwestern Nigeria. *J Contemp Med Sci*. 2018;4.
- Eftekhari Vaghefi SH, Sheikhabaehi F, Mokhtari T, Khademi F, Bahari H, Ghorbani R. A model for individual height estimation from forearm length in natives of Kerman, Iran. *Anat Sci J*. 2014;11:141–144.
- Moshkdanian G, Mahaki Zadeh S, Moghani Ghoroghi F, Mokhtari T, Hassanzadeh G. Estimation of Stature from the Anthropometric Measurement of Lower Limb in Iranian Adults. *Anat Sci J*. 2014;11:149–154.
- Tsokos M, Türk EE, Madea B, Koops E, Longauer F, Szabo M, et al. Pathologic features of suicidal deaths caused by explosives. *Am J Forensic Med Pathol*. 2003;24:55–63.
- Krishan K, Sharma A. Estimation of stature from dimensions of hands and feet in a North Indian population. *J Forensic Leg Med*. 2007;14(6):327–332.
- Meadows L, Jantz RL. Estimation of stature from metacarpal lengths. *J Forensic Sci*. 1992;37:147–154.
- Jasuja OP, Singh G. Estimation of stature from hand and phalange length. *J Indian Acad Forensic Med*. 2004;26:0971–0973.
- Zeybek G, Ergur I, Demiroglu Z. Stature and gender estimation using foot measurements. *Forensic Sci Int*. 2008;181:54. e1–e5.
- Ruff CB, Holt BM, Niskanen M, Sladěk V, Berner M, Garofalo E, et al. Stature and body mass estimation from skeletal remains in the European Holocene. *Am J Phys Anthropol*. 2012;148:601–617.
- Spradley MK, Jantz RL. Sex estimation in forensic anthropology: skull versus postcranial elements. *J Forensic Sci*. 2011;56:289–296.
- Celbis O, Agritmis H. Estimation of stature and determination of sex from radial and ulnar bone lengths in a Turkish corpse sample. *Forensic Sci Int*. 2006;158:135–139.
- Sanli SG, Kizilkanat ED, Boyan N, Ozsahin ET, Bozkir MG, Soames R, et al. Stature estimation based on hand length and foot length. *Clin Anat*. 2005;18:589–596.
- Kanchan T, Menezes RG, Moudgil R, Kaur R, Kotian MS, Garg RK. Stature estimation from foot dimensions. *Forensic Sci Int*. 2008;179:241. e1–241.e5.
- Evereklioglu C, Doganay S, Er H, Gunduz A, Tercan M, Balat A, et al. Craniofacial anthropometry in a Turkish population. *Cleft Palate Craniofac J*. 2002;39:208–218.
- Kau CH, Richmond S, Incrapera A, English J, Xia JJ. Three-dimensional surface acquisition systems for the study of facial morphology and their application to maxillofacial surgery. *Int J Med Robot*. 2007;3:97–110.
- de Acuerdo PdE, de la Cabeza TM. Prediction of Stature According to Three Head Measurements. *Int J Morphol*. 2015;33:1151–1155.
- Krishan K, Kumar R. Determination of stature from cephalo-facial dimensions in a North Indian population. *Leg Med (Tokyo)*. 2007;9:128–133.
- Sahni D, Sanjeev, Sharma P, Harjeet, Kaur G, Aggarwal A. Estimation of stature from facial measurements in northwest Indians. *Leg Med (Tokyo)*. 2010;12:23–27.
- Ibrahim MA, Khalifa AM, Hassan HA, Tamam HG, Hagrais AM. Estimation of stature from hand dimensions in North Saudi population, medicolegal view. *Saudi J Forensic Med Sci*. 2018;1:19–27.
- Kumari KL, Vijaya Babu PVSS, Kusuma Kumari P, Nagamani M. A study of cephalic index and facial index in Visakhapatnam, Andhra Pradesh, India. *Int J Res Med Sci*. 2015;3:656–658.
- Poorhassan M, Mokhtari T, Navid S, Rezaei M, Sheikhezadi A, Mojavverostami S, et al. Stature estimation from forearm length: an

- anthropological study in Iranian medical students. *J Contemp Med Sci*. 2017;3:270–272.
22. Mahakizadeh S, Moghani-Ghoroghi F, Moshkdanian Gh, Mokhtari T, Hassanzadeh G. The determination of correlation between stature and upper limb and hand measurements in Iranian adults. *Forensic Sci Int*. 2016;260:27–30.
 23. Navid S, Mokhtari T, Alizamir T, Arabkheradmand A, Hassanzadeh G. Determination of stature from upper arm length in medical students. *Anat Sci J*. 2014;11:135–140.
 24. Michopoulou E, Ekizoglu O, Er A, Bozdog M, Akcaoglu M, Can IO, et al. Stature estimation using long bone metrics in contemporary Turkish. *La Revue de Médecine Légale*. 2017;8:190.
 25. Din TNDT, Rajion ZA, Luddin N. Nasofacial morphometric analysis for nasal reconstruction. *J Teknol*. 2015;76:81–85.
 26. Omotoso DR, Oludiran OO, Sakpa CL. Nasofacial Anthropometry of Adult Bini Tribe in Nigeria. *Afr J Biomed Res*. 2011;14:219–221.
 27. Krishan K. Estimation of stature from cephalo-facial anthropometry in north Indian population. *Forensic Sci Int*. 2008;181:52. e1–e6.
 28. Chiba M, Terazawa K. Estimation of stature from somatometry of skull. *Forensic Sci Int*. 1998;97:87–92.
 29. Kalia S, Shetty SK, Patil K, Mahima VG. Stature estimation using odontometry and skull anthropometry. *Indian J Dent Res*. 2008;19:150–154.
 30. Sarangi AK, Dadhi B, Mishra K. Estimating of stature from adult skull bone. *J Ind Acad Forensic Med*. 1981;182:24–26.
 31. Agnihotri AK, Kachhwaha S, Googoolye K, Allock A. Estimation of stature from cephalo-facial dimensions by regression analysis in Indo-Mauritian population. *J Forensic Leg Med*. 2011;18:167–172.

This work is licensed under a Creative Commons Attribution-NonCommercial 3.0 Unported License which allows users to read, copy, distribute and make derivative works for non-commercial purposes from the material, as long as the author of the original work is cited properly.